

RESEARCH ARTICLE

Open Access

Time-course microarrays reveal early activation of the immune transcriptome and adipokine dysregulation leads to fibrosis in visceral adipose depots during diet-induced obesity

Eun-Young Kwon¹, Su-Kyung Shin¹, Yun-Young Cho¹, Un Ju Jung¹, Eunjung Kim², Taesun Park³, Jung Han Yoon Park⁴, Jong Won Yun⁵, Robin A McGregor¹, Yong Bok Park⁶ and Myung-Sook Choi^{1,7*}

Abstract

Background: Visceral white adipose tissue (WAT) hypertrophy, adipokine production, inflammation and fibrosis are strongly associated with obesity, but the time-course of these changes *in-vivo* are not fully understood. Therefore, the aim of this study was to establish the time-course of changes in adipocyte morphology, adipokines and the global transcriptional landscape in visceral WAT during the development of diet-induced obesity.

Results: C57BL/6 J mice were fed a high-fat diet (HFD) or normal diet (ND) and sacrificed at 8 time-points over 24 weeks. Excessive fat accumulation was evident in visceral WAT depots (Epididymal, Perirenal, Retroperitoneum, Mesentery) after 2–4 weeks. Fibrillar collagen accumulation was evident in epididymal adipocytes at 24 weeks. Plasma adipokines, leptin, resistin and adiponectin, increased early and time-dependently, while adiponectin decreased late after 20 weeks. Only plasma leptin and adiponectin levels were associated with their respective mRNA levels in visceral WAT. Time-course microarrays revealed early and sustained activation of the immune transcriptome in epididymal and mesenteric depots. Up-regulated inflammatory genes included pro-inflammatory cytokines, chemokines (Tnf, Il1rn, Saa3, Emr1, Adam8, Itgam, Ccl2, 3, 4, 6, 7 and 9) and their upstream signalling pathway genes (multiple Toll-like receptors, Irf5 and Cd14). Early changes also occurred in fibrosis, extracellular matrix, collagen and cathepsin related-genes, but histological fibrosis was only visible in the later stages.

Conclusions: In diet-induced obesity, early activation of TLR-mediated inflammatory signalling cascades by CD antigen genes, leads to increased expression of pro-inflammatory cytokines and chemokines, resulting in chronic low-grade inflammation. Early changes in collagen genes may trigger the accumulation of ECM components, promoting fibrosis in the later stages of diet-induced obesity. New therapeutic approaches targeting visceral adipose tissue genes altered early by HFD feeding may help ameliorate the deleterious effects of diet-induced obesity.

Keywords: Adipocytokine dysregulation, Transcriptional response, Adipose tissue, Extracellular matrix, Cathepsin, Fibrosis

* Correspondence: mschoi@knu.ac.kr

¹Center for Food and Nutritional Genomics Research, Kyungpook National University, Daegu, Republic of Korea

⁷Department of Food Science and Nutrition, Kyungpook National University, 1370 Sank-Yuk Dong Puk-Ku, Daegu 702-701, Korea

Full list of author information is available at the end of the article

Background

White adipose tissue (WAT) not only serves as a fuel storage depot but also releases a variety of bioactive factors, called adipokines, which have pro- and anti-inflammatory activity. In obesity, most adipokines, including leptin, resistin and tumor necrosis factor alpha (TNF α), are overproduced and promote inflammation. Whereas, the production of anti-inflammatory adipokines, such as adiponectin, is decreased with increasing adiposity. Since, inflammation is considered to be causal in the development of obesity and insulin resistance [1-3], the dysregulated production of adipokines caused by excessive adipose tissue expansion is linked to the pathophysiology of obesity-related metabolic syndrome [4].

There is a close relationship between inflammation and the extracellular matrix (ECM), which requires remodelling to accommodate adipocyte growth during adipose tissue expansion. In obesity, adipose tissue inflammation is further exacerbated by excessive accumulation of ECM components in adipose tissue, possibly as a result of pro-inflammatory cytokines released by adipocytes [5]. Excessive accumulation of ECM components may represent a key event in obesity-mediated adipose tissue fibrosis, similar to fibrosis in other tissues such as liver [6,7]. Accumulating evidence indicates that altered circulating adipokine levels contribute to the progression of cardiac or hepatic fibrosis [8,9]. Thus, adipokine dysregulation may also regulate remodeling of the adipocyte ECM and fibrosis. However, the actual time-course of adipokine dysregulation, inflammation, ECM accumulation leading to the development of adipose tissue fibrosis is yet to be fully established.

High-throughput microarrays assessing global gene expression changes provide a unique window into the molecular events occurring in adipose tissue during the development of obesity. Previous microarray studies have revealed that a high fat diet (HFD) is associated with adipokine dysregulation, inflammation and fibrosis [10-15]. However, these microarray studies tend to compare gene expression differences at a single end-point only or over short-time periods between 2–12 weeks. Following the molecular changes in adipose tissue during the development of obesity in humans is not feasible. Therefore, time-course models of diet-induced obesity in mice can help improve our understanding of the molecular pathogenesis of obesity *in-vivo*. However, the use of genetically obese animals or extremely high levels of dietary fat intake, which do not reflect normal dietary consumption hampers the translation of findings from mouse obesity models to humans [10-14]. Therefore we established a long-term diet-induced obesity model, by feeding C57BL/6 J mice with a high-fat diet (39% kcal) or normal diet for 24 weeks and sacrificing mice at regular time-points to allow in-depth time-course microarray analysis of visceral adipose tissue depots.

Accordingly the aim of the present study was to establish the time-course of changes in adipocyte morphology, adipokines and the global transcriptional landscape in visceral adipose tissue depots during the transition from a lean to obese phenotype in HFD fed mice. We primarily focus on the timing of changes associated with adipokine levels, inflammation and fibrosis, which are recognized as important factors associated with human obesity.

Methods

Animals

Three hundred and sixty male inbred C57BL/6 J mice were obtained from the Jackson Laboratory (Bar Harbor, ME) at 4 weeks of age. All the mice were individually housed under a constant temperature (24°C) and 12-hour light/dark cycle, fed the AIN-76 semi-purified diet for one week acclimation period after arrival, and then randomly divided into a normal diet (ND) and an HFD group, with 18 mice per group for 0, 2, 4, 6, 8, 12, 16, 20 and 24 weeks. After the acclimation, the HFD group was fed the AIN-76 semi-purified diet with 39.2 kcal% fat (American Institute of Nutrition, 1977) consisting of 20% fat and 1% cholesterol (Additional file 1: Table S1). Every 2–4 weeks, after a 12 h fast, blood samples were drawn from the inferior vena cava. Blood was collected in heparin-coated tubes, centrifuged at 1,000 \times g for 15 min at 4°C and then stored at -70°C. Mice were anaesthetized and sacrificed at 0, 2, 4, 6, 8, 12, 16, 20 and 24 weeks. The adipose tissues were then removed, rinsed, weighed, immediately frozen in liquid nitrogen, and stored at -70°C until further analysis. The current study protocol was approved by the Ethics Committee for animal studies at Kyungpook National University, Republic of Korea.

Plasma adipokines

Radioimmunoassay were performed for the measurement of plasma amylin (amylin RIA kit; Diagnostic Systems Laboratories), adiponectin (adiponectin RIA kit; PerkinElmer Life and Analytical Sciences, Boston, MA), leptin (Mouse leptin RIA kit; Linco Research, St. Charles, MO), and resistin concentrations (resistin RIA kit; PerkinElmer Life and Analytical Sciences, Boston, MA).

Morphology of fat tissues

Epididymal adipose tissue was fixed in 10% (v/v) paraformaldehyde/PBS and embedded in paraffin for staining with hematoxylin and eosin (H&E) and Masson's trichrome. The stained area was viewed using a microscope at a magnification of 200 \times and 400 \times , respectively.

RNA preparation and quality control

Total RNA was extracted from the adipose tissue using TRIZOL reagent (Invitrogen Life Technologies, Grand Island, NY) according to the manufacturer's instructions. DNase digestion was used to remove any DNA contamination and RNA was re-precipitated in ethanol to ensure no phenol contamination. For quality control, RNA purity and integrity were evaluated using the Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, USA). For epididymal adipose tissue, six pooled RNA sample sets were constructed for the ND and HFD groups at each time-point as described previously [16]. In mesenteric adipose tissue, one pooled RNA sample set was constructed for the ND and HFD groups at each time-point based on 18 individual samples. RNA was stored at -70°C prior to further analysis by microarray and RT-qPCR.

Microarray analysis

Total RNA was amplified and purified using the Ambion Illumina RNA amplification kit (Ambion, Austin, USA) to yield biotinylated cRNA according to the manufacturer's instructions. After purification, the cRNA was quantified using the ND-1000 Spectrophotometer (NanoDrop, Wilmington, USA). 750 ng of labelled cRNA samples were hybridized to each Illumina MouseWG-6 v2 Expression BeadChip for 16–18 h at 58°C , according to the manufacturer's instructions (Illumina, Inc., San Diego, USA). Detection of array signal was carried out using Amersham fluorolink streptavidin-Cy3 (GE Healthcare Bio-Sciences, Little Chalfont, UK) following the bead array manual. The quality of hybridization and overall chip performance were monitored by visual inspection of both internal quality controls and the raw scanned data. Raw data was extracted using the Illumina BeadStudio software provided by the manufacturer. Probe signal intensities significantly higher than background intensities were determined (detection p -value > 0.05), but probe signal data were not filtered to preserve probes with low expression at different time points. Probe signal intensities were quantile normalized and log transformed. Microarray analysis was performed in ArrayAssist (Stratagene, USA), Bioconductor and R programming language. LIMMA was used to determine significant differentially expressed genes (HFD responsive genes) between HFD- and ND-fed mice at each time point based on FDR $< 5\%$, Benjamin and Hochberg adjusted p -value < 0.05 and log fold change > 1 [17]. This microarray data was deposited in Gene Expression Omnibus (GEO) database (GEO accession numbers: GSE39549).

RT-qPCR

The same total RNA (1 μg) samples that were hybridised to microarrays were reverse-transcribed into cDNA

using the QuantiTect[®] reverse transcription kit (Qiagen, Germany). Then mRNA expression was quantified by real-time quantitative PCR, using the SYBR green PCR kit (Qiagen, Germany) and the CFX96TM real-time system (BIO-RAD). Gene specific mouse primers were used to detect leptin, adiponectin and resistin (Additional file 2: Table S2). Cycle thresholds were determined based on SYBR green emission intensity during the exponential phase. Ct data were normalized using Gapdh, which was stably expressed in both HFD and ND fed mice. Relative gene expression was calculated using the $2^{-\Delta\Delta\text{Ct}}$ method [18].

Functional features of high-fat feeding responsive genes in visceral adipose tissues

To identify which biological processes were associated with the visceral adipose tissue genes altered by HFD at one or more time-points over 24 weeks, enriched functional features were identified using DAVID [19]. The analysis was performed using Gene Ontology and Pathways and mapped by DAVID ID. Then the DAVID Functional Annotation Clustering tool [19] was applied to identify enriched biological themes and cluster redundant annotation terms.

Identification of genes that respond to both age and high-fat diet simultaneously in epididymal adipose tissue

Next, we examined which HFD-responsive genes in epididymal adipose tissue were correlated with age and/or body mass. At first, genes affected by advancing age were identified by examining genes that were continuously up- or down-regulated in the both ND and HFD group. These genes were overlaid with the HFD-responsive genes (HFD vs. ND, fold change > 2 and p -value < 0.05) to screen genes that responded to both age and high-fat diet simultaneously in epididymal adipose tissue. These overlapped genes were selected and correlated with body mass.

Statistical analysis

The parameter values were expressed as the mean (standard error of the mean (SEM)). Significant differences between groups were determined by student's t -test and wilcoxon t -test using the SPSS program (SPSS Inc., Chicago, IL). Correlation analyses utilized the Pearson's coefficient. Results were considered statistically significant at $p < 0.05$.

Results

Time-course of changes in body weight and food efficiency during diet-induced obesity

Significant increases of body weight gain and energy consumption were observed in the HFD group after the 4th week and 2th week, respectively. The food efficiency

ratio (FER) was also significantly increased in the HFD group after the 4th week of high-fat feeding (Additional file 3: Table S3) [20] compared to the ND group.

Time-course of changes in WAT depot weights, adipocyte size and adipose tissue fibrosis during diet-induced obesity

When WAT weights were expressed in g/100 g WAT, visceral WAT depots including epididymal, perirenal, retroperitoneum, mesentery WAT were significantly increased from week 2 in the HFD group compared to ND group. Similarly, from week 2, subcutaneous WAT was significantly higher in the HFD group (Figure 1). In addition to increased body fat, adipocyte size was also enlarged in the epididymal WAT depot in a time-dependent manner (Figure 2A).

To assess fibrosis Masson's trichrome staining was performed, which revealed the epididymal WAT depot of the HFD group contained very pronounced trichrome-positive "streaks" interspersed among the adipocytes at 24 weeks. Whereas, the epididymal WAT depot of the ND group revealed a normal fat-pad, with densely packed and very thin collagen sheets surrounding each adipocyte at week 24 (Figure 2B).

Time-course of changes in plasma adipokine, glucose, insulin and lipids levels during diet-induced obesity

The time-course of changes in plasma adipokine levels in response to HFD is shown in Figure 3. The most prominent changes were observed in leptin and resistin levels, which were increased from week 4 to week 24 in the HFD group. Plasma adiponectin levels were also elevated in the HFD group from week 4, reaching a peak at week 6 and thereafter gradually decreasing. In contrast, plasma adiponectin levels in the HFD group tended to be lower than in the ND group for the overall feeding period and were significantly lower at week 20 and 24. Fasting blood glucose and insulin resistance and plasma total cholesterol concentration were markedly elevated in the HFD group after 16 weeks and 6 weeks, respectively. No differences in plasma TG and FFA levels were detectable between the HFD group and ND group at any time points (data not shown) [21].

Time-course of transcriptional changes in epididymal and mesenteric WAT depots during diet-induced obesity

In the epididymal WAT depot, 2037 genes were altered at one or more time points in response to HFD intake over 24 weeks (Table 1), of which 1610 genes were up-regulated and 445 genes were down-regulated. In mesenteric WAT depot, 1784 genes were altered at one or more time point in response to HFD intake over 24 weeks, of which 1134 genes were up-regulated and 892 genes were down-regulated (Table 1). Functional

gene ontology terms associated with these HFD responsive genes in the epididymal WAT and mesenteric WAT depots were clustered using DAVID (Table 2). Immune and inflammatory system function terms were enriched among the consistently up-regulated genes across all time-points, in response to HFD, in both the epididymal and mesenteric WAT depots (Table 2A). Glucose metabolic system function terms were enriched among the down-regulated genes (Table 2B). Lipid and cholesterol metabolism related functional gene ontology terms were specifically enriched among the down-regulated genes in the mesenteric WAT depot across all time-points.

Epididymal WAT depot genes altered by advancing age and diet-induced obesity

When HFD-responsive genes, in epididymal WAT, were overlapped with age-dependent genes, 14 genes were identified as shown in Figure 4. Fold changes and q-values of identified 14 genes are given in Additional file 4: Table S4. These 14 genes were correlated with body weight in both ND and HFD groups. Accordingly, these genes were associated with not only with age but also diet-induced obesity. In particular, among these 14 age and HFD responsive genes was EGF-like module containing, mucin-like, hormone receptor-like sequence 1 (*Emr1*) and chemokine (C-C motif) ligand 7 (*Ccl7*), which are both genes associated with inflammation, as well as body weight gain.

Time-course of transcriptional changes in adipokine, pro-inflammatory cytokine, chemokine and upstream signalling genes in visceral WAT depots during diet-induced obesity

The time-course of pro- and anti-inflammatory adipokine gene expression changes in WAT is shown in Figure 5. Consistent with plasma leptin and adiponectin levels, leptin mRNA expression was also up-regulated, while adiponectin mRNA expression was down-regulated consistently during the development of diet-induced obesity over 24 weeks. Microarray validation using RT-qPCR revealed agreement in the time-course and pattern of changes in gene expression. There was no significant change in resistin mRNA expression.

During the development of diet-induced obesity, transcription of several pro-inflammatory cytokines and chemokines such as disintegrin and metalloproteinase domain 8 (*Adam8*), *Emr1*, interleukin 1 receptor antagonist (*Il1rn*), tumor necrosis factor (*Tnf*), serum amyloid A 3 (*Saa3*), integrin alpha-M (*Itgam*), chemokine (C-C motif) ligand 2 (*Ccl2*), ligand 3 (*Ccl3*), ligand 4 (*Ccl4*), ligand 6 (*Ccl6*), ligand 7 (*Ccl7*) and ligand 9 (*Ccl9*), were significantly elevated in epididymal WAT between week 20 and 24 (Fig. 6A). Moreover, gene expression of toll-like receptors (TLRs), interferon regulatory factor 5




Figure 1 (See legend on next page.)

(See figure on previous page.)

Figure 1 Timely effect of high-fat feeding on adipose tissue weights over 24 weeks in C57BL/6 J mice. Data shown as means \pm S.D. Values are significantly different from the ND group according to Student's test: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. ND: normal diet (AIN-76), HFD: high-fat diet (20% fat, 1% cholesterol), WAT: white adipose tissue.

(*Irf5*) and CD antigen families were also significantly altered by HFD in epididymal WAT. Among them *Tlr2*, *Tlr6*, *Tlr7*, *Tlr8*, *Tlr13* and *Trl1*, *Irf5* were significantly up-regulated by HFD from week 8 and week 20, respectively (Figure 6B, C). Expression of *Cd9*, *Cd14*, *Cd37*, *Cd44*, *Cd68*, *Cd72*, *Cd83*, *Cd84*, *Cd180*, *Cd276*, *Cd300a* and *Cd300b* was also significantly elevated from week 8 to 24 in a time dependent manner (Figure 6D).

Time-course of transcriptional changes in extracellular matrix genes, adipose tissue-expressed collagen genes and cathepsin family genes during diet-induced obesity

To identify the time-dependent effect of HFD on fibrosis relative to the ND group, we examined the levels of many key fibrotic genes (Figure 7). Lumican (*Lum*) and transforming growth factor β (*Tgfb1*) mRNA levels were up-regulated, while decorin (*Dcn*) mRNA levels were down-regulated by HFD in a time dependent manner. Several matrix metalloproteases (*Mmp*) showed time-dependent transcriptional alterations in response to HFD. *Mmp2*, *Mmp3* and *Mmp13* mRNA levels were up-regulated by HFD from week 8 to 24, whereas *Mmp12* mRNA levels were significantly up-regulated earlier after week 2 in the HFD group (Figure 7A). Collagen-related mRNA levels including *Col1a2*, *Col3a1*, *Col4a5*, *Col8a1*, *Col9a3* and *Col16a1* were significantly higher at week 2 in the HFD group, relative to the ND group, and sharply decrease after 4 weeks (Figure 7B). Cathepsin (*Cts*) mRNA levels were also altered by HFD (Figure 7C), *Ctsa*, *Ctsb*, *Ctsk*, *Ctsl*, *Ctss* and *Ctsz* mRNA levels were mostly up-regulated by high-fat feeding in the epididymal WAT depot. In particular, *Ctsk* mRNA levels were significantly increased from week 8 in the HFD group. *Ctss* and *Ctsz* mRNA levels were significantly higher at week 2, 8, 20 and 24, while *Ctsb* mRNA levels were only higher at week 2 and week 24 in the HFD group.

Discussion

Adipose tissue can rapidly expand in response to excess energy intake through both adipocyte hypertrophy (cell size increase) and hyperplasia (cell number increase). Excessive adipose tissue expansion in obesity is associated with adipokine dysregulation, inflammation, ECM remodeling and fibrosis particularly in visceral adipose depots. However, the time-course of these pathophysiological events and the underlying transcriptional changes in adipose tissue are not fully understood. We show adipose tissue expansion occurs in multiple visceral WAT

depots early during the development of diet-induced obesity. Pro-inflammatory adipokine levels also increase early after 2–4 weeks, while anti-inflammatory adipokine levels are reduced after 20 weeks in HFD fed mice. Time-course microarray profiling revealed early and prolonged up-regulation of inflammation and immune system associated genes in the epididymal WAT depot and the smaller mesenteric WAT depot. In the later stages of diet-induced obesity, fibrosis was evident in the epididymal WAT depot. Hence the long-term diet-induced obesity model used in the present study provides a unique insight into the timing of events in visceral adipose tissue during the development of obesity.

Adipocytokine dysregulation is reported to be closely associated with obesity and plasma proinflammatory adipokines are widely reported to be elevated in obese animals. Adipokines are secreted from visceral adipose depots, hence expansion of visceral adipose tissue depots is correlated with plasma adipokine levels, but it is less well established whether adipokine are transcriptionally regulated in obesity. In the present study, plasma leptin and resistin levels were increased consistently from 4 to 24 weeks. Time-course transcriptional profiling revealed leptin mRNA levels were increased in both the epididymal and mesenteric WAT depots over 24 weeks, but changes in resistin mRNA levels were less clear. Notably, leptin mRNA levels were higher in the mesenteric WAT depot than in the epididymal WAT depot. From a structural perspective, leptin is a member of the cytokine family, and leptin receptors belong to the class-I cytokine receptor family [22,23]. Leptin can control the production and activation of pro-inflammatory cytokines such as IL-6 and TNF- α by macrophages [24,25], and is a key regulatory factor expressed in both zebrafish and mammalian obesity in lipid metabolism [26]. Resistin is also known as an adipose tissue-specific secretory factor, participating in the pathogenesis of insulin resistance, adipogenesis and inflammation in mice [27,28]. However, the time-course of plasma resistin level changes, was largely dissociated from resistin mRNA levels in the epididymal and mesenteric WAT depots. Jackson et al. [29] reported similar observations, largely that resistin mRNA levels were very low or undetectable in adipose tissue, which suggests that visceral adipocytes are not a major cell type producing resistin. Plasma resistin levels may be due to resistin production by circulating monocytes and macrophages. Like many other pro-inflammatory cytokines, resistin stimulates intracellular





Figure 2 Timely effect of high-fat feeding over 24 weeks on (A) H&E and (B) Masson's trichrome staining of epididymal white adipose tissue in C57BL/6 J mice. (A) H&E stained transverse-section of epididymal fat. Original magnification $\times 200$. (B) Masson's trichrome stained transverse-section of epididymal fat. Fibrillar collagens, primarily collagen I and III, are stained with blue as indicated with arrowheads. Nuclei and keratin are stained with deep purple and red, respectively. Original magnification $\times 400$. ND: normal diet (AIN-76), HFD: high-fat diet (20% fat, 1% cholesterol).

signalling through NF- κ B activation, which in turn promotes the synthesis of other pro-inflammatory cytokines, including MCP-1, vascular cell adhesion molecule-1 (VCAM-1), as well as intercellular adhesion molecule-1 (ICAM-1), which are involved in chemotactic pathways directing leukocyte recruitment to sites of infection [30,31]. Therefore, while resistin appears not to be produced by visceral adipose tissue depots, plasma resistin may still be a player in early visceral adipose tissue inflammation via activation of pro-inflammatory signalling and leukocyte recruitment.

Circulating plasma adiponectin and adiponectin mRNA levels are reportedly decreased by HFD and inversely correlated with fat mass, and directly correlated with insulin sensitivity [32,33]. In the present study, a markedly different time-course in plasma adiponectin levels was evident with late decreases in plasma adiponectin observed, along with concomitant down-regulation of adiponectin mRNA levels in both the epididymal and mesenteric WAT depots. There have been reports indicating adiponectin is

probably not a useful early biomarker of obesity. Sumiyoshi and colleagues reported that high fat feeding (45% fat, wt/wt) for 55 weeks did not alter circulating adiponectin level in C57BL/6 J mice compared with a low fat diet (3% fat, wt/wt) [34]. In accordance with our results, Park and colleague's reported that circulating adiponectin level was significantly reduced after 20 weeks of high-fat feeding (55 kcal% from fat) compared with a normal diet [35]. The lack of any early changes in plasma adiponectin and visceral adipose tissue adiponectin mRNA levels in HFD fed mice may be part of an adaptive response to protect against early inflammatory changes. There were prominent changes in adiponectin levels in both HFD and ND fed mice around 6 weeks, which we speculate may be related to sexual maturation or ageing.

Adipsin, a serine protease is synthesized and secreted by adipose cells and is found in the bloodstream [36]. In the current study, plasma adipsin levels were significantly higher in the HFD group compared to ND group from the week 2 to 24. However, it is not clear whether



the higher concentration of adipisin in blood reflects increased adiposity in the HFD fed mice, because fat mass was increased over time but plasma adipisin decreased after 6 weeks regardless of diet. Previous

Table 1 Number of high-fat diet responsive genes in two visceral WAT depots at multiple time-points in C67BL/6 J mice

Time (week)	Epididymal WAT			Mesenteric WAT		
	Fold-changed genes	▲	▼	Fold-changed genes	▲	▼
2	1478	1233	245	999	684	315
4	3	1	2	190	177	13
8	37	34	3	306	49	257
20	339	200	139	247	32	215
24	431	341	90	751	445	306
All	2037	1610	445	1784	1134	892

Differentially expressed genes based on HFD v ND comparison at each time-point according to p -value < 0.05 , fold change > 2 in the epididymal WAT depot and fold change > 2 in the mesenteric WAT depot. Unique HFD responsive genes were differentially expressed at one or more time-points. ▲: up-regulated, ▼: down-regulated, WAT: white adipose tissue.

studies have reported that blood adipisin levels are substantially increased in obese humans [37,38], but decreased in murine models of obesity [39]. Taken together, the present *in-vivo* observations of the time-course of various plasma adipokine levels and visceral WAT depot adipokine mRNA levels suggests leptin gene expression is up-regulated early, while adiponectin gene expression is down-regulated later in the epididymal and mesenteric WAT depots contributing to time-dependent changes in plasma adipokine levels.

We conducted in-depth time-course microarray profiling. The early and prolonged changes in the global transcriptome of both epididymal and mesenteric WAT revealed up-regulation of genes predominantly associated with immune and inflammatory system function enriched across all time-points in HFD fed mice. On the other hand, down-regulated genes were predominantly associated with monosaccharide, hexose and glucose metabolism across all time-points. Interestingly, down-regulated genes were also associated with lipid and cholesterol metabolism across all time-points although predominantly in the mesenteric WAT depot, rather than the epididymal

Table 2 Functional annotation gene clusters (A) up-regulated and (B) down-regulated in visceral adipose tissue depots by high-fat feeding in C57BL/6 J mice over 24 weeks


	Depot	E.S.	Functional terms	No. of genes	p-value		
(A)	Epididymal WAT	5.34	Inflammatory response	41	1.10E-06		
			Response to wounding	53	6.40E-06		
			Defense response	62	8.20E-06		
		Mesenteric WAT	4.84	2.59	Regulation of phagocytosis	10	4.20E-05
				2.23	Cell activation	38	5.60E-05
				2.09	Positive regulation of immune system process	37	2.30E-06
					Positive regulation of immune response	25	8.40E-05
	Mesenteric WAT	3.48		Positive regulation of immune system process	32	3.80E-08	
				Positive regulation of immune response	25	5.40E-08	
				Activation of immune response	17	4.40E-06	
				Defense response	47	3.40E-06	
			2.72	Cell activation	32	2.10E-06	
				Leukocyte activation	28	1.50E-05	
				Lymphocyte activation	25	3.10E-05	
(B)	Epididymal WAT	5.77	Fatty acid metabolic process	20	1.60E-08		
			3.55	Hexose metabolic process	20	2.80E-09	
				Monosaccharide metabolic process	21	3.40E-09	
				Glucose metabolic process	17	3.40E-08	
	Mesenteric WAT	12.62		Glucose metabolic process	36	2.80E-18	
				Monosaccharide metabolic process	40	8.20E-17	
				Hexose metabolic process	37	4.80E-16	
				Glycolysis	17	7.60E-12	
				Gluconeogenesis	7	9.10E-05	
			4.41	Steroid metabolic process	26	3.80E-08	
				Sterol metabolic process	16	9.10E-07	
				Cholesterol metabolic process	15	1.50E-06	
				Lipid biosynthetic process	32	1.90E-06	
				Lipid catabolic process	20	5.60E-06	
	Acetyl-CoA metabolic process	9	3.70E-05				

Functional annotation terms enriched amongst the up/down regulated genes in epididymal and mesenteric WAT of HFD fed C57BL/6 J mice over 24 weeks. Functional annotation terms were clustered according to biological processes. WAT: white adipose tissue; E.S.: Enrichment Score.

WAT depot. In agreement with other diet-induced obesity models, functional ontology terms including sterol metabolism and acetyl-CoA biosynthesis were reportedly highly enriched among regulated genes in the mesenteric WAT depot but not in the epididymal WAT depot [40], and transcripts involved in lipid biosynthesis were up-regulated only by low fat meals [33]. Genes associated with the inflammatory response that were significantly up-regulated in the epididymal WAT depot during diet-induced obesity included pro-inflammatory cytokine and chemokine genes (*Tnf*, *Il1rn*, *Saa3*, *Emr1*, *Adam8*, *Itgam*, *Ccl2*, *Ccl3*, *Ccl4*, *Ccl6*, *Ccl7* and *Ccl9*) and their upstream signalling pathway genes such as TLRs, *Irf5* and *Cd14*. In accordance with our study, many inflammatory genes

including *Ccl3*, *Ccl2*, *Saa3*, *Tlr13*, which were highly expressed in adipose tissue of fat mouse, have been linked with obesity and/or insulin resistance [41], furthermore *Emr1* and *Ccl7* mRNA expression are associated with body weight gain as well as obesity-related inflammation. Adipose tissue consists of multiple cell types including adipocytes, fibroblasts, macrophages and endothelial cells. Therefore, the genes identified in the present study as differentially expressed during the development of diet-induced obesity will reflect overall changes in visceral adipose tissue depots rather than specific cell-types.

Recent studies have suggested some interplay between TLRs and adipokines. Lin et al. [42] suggested adipocyte *Tlr2* synthesis increases upon stimulation



with *Tnf*. Also, the expression of *Tlr1-9* and *Tlr11-13* in murine adipose tissue appears to be strongly regulated by leptin [43]. In the present study, the expression of *Tlr1*, *Tlr2*, *Tlr6*, *Tlr7*, *Tlr8* and *Tlr13* were significantly up-regulated by HFD in the epididymal WAT depot, along with *Tnf* and *Lep* expression, while expression of *Adipoq* was down-regulated. HFD also significantly up-regulated *Irf5* in the later stages of diet-induced obesity. *Irf5* is generally involved downstream of the TLR-MyD88 signalling pathway for induction of pro-inflammatory cytokines, and *Irf5* is a potential target of therapeutic interventions to control harmful immune responses [44].

Of the thirteen TLRs, TLR4 was characterized first, working in conjunction with the LPS receptor CD14, TLR4 is the primary signalling receptor for gram negative bacterial

lipopolysaccharide (LPS) [45]. CD14 is crucial for LPS recognition by TLR4 and cooperates with other TLRs, including TLR2 and TLR3 [46,47]. Cd180 also belongs to the family of pathogen receptors. Cd180 takes part in B-cell recognition and LPS signalling [48]. In this study, we show diet-induced obesity leads to significant induction of *Cd14*, *Cd180*, *Cd68* and other Cd antigen genes active in the immune and inflammation system of animals. Therefore, expression of Cd antigen genes may be used as markers of visceral adipose tissue inflammation during the development of diet-induced obesity. Taken together these findings suggest that, adipose tissue expansion, and concomitant activation of TLR-mediated inflammatory signalling cascades and induction of CD antigens, causes increased expression of pro-inflammatory cytokines and chemokines, which results in chronic low-grade inflammation.




Figure 5 RT-qPCR validation. Time-dependent fold changes in (A) Lep mRNA, (B) Adipoq mRNA and (C) Retn mRNA expression based on HFD vs. ND group over 24 weeks. Microarray and RT-qPCR data shown as means \pm S.D. * $p < 0.05$ based on wilcoxon t-test and student's t-test, respectively. No statistical analysis, since we evaluated 1 pooled sample for microarray analysis in MAT. ND: normal diet (AIN-76), HFD: high-fat diet (20% fat, 1% cholesterol), EAT: epididymal adipose tissue, MAT: mesenteric adipose tissue, Lep: leptin, Adipoq: adiponectin, Retn: resistin.

Obesity is in most, but not all humans, associated with marked changes in the secretory function of adipocytes and macrophages, together with chronic low-grade inflammation and an increased risk of developing fibrosis. A complex network of inflammatory cytokines, adipokines, transcription factors, and receptor molecules is involved in the development of adipose tissue fibrosis. In

adipose tissue, fibrosis appears to be initiated in response to adipocyte hypertrophy. Adipose tissue ECM remodelling, which plays a pivotal role in adipogenesis and tissue architecture [49], is crucial to accommodate obesity-induced cellular alterations [50]. However, the persistence of an inflammatory stimulus in adipose tissue may be responsible for the excessive synthesis of





Figure 6 Effect of high-fat feeding on transcription of (A) pro-inflammatory cytokine and chemokine genes, (B) Toll-like receptors, (C) IRF5 and (D) CD families expressed in epididymal adipose tissue of C57BL/6 J mice over 24 weeks (HFD vs. ND group). mRNA expression in epididymal adipose tissue measured by the microarray analysis. Fold changes in mRNA expression (HFD vs. ND) were displayed at each time point over 24 weeks. Data shown as means \pm S.D. * $p < 0.05$ based on wilcoxon t-test. ND: normal diet. HFD: high-fat diet.

ECM components and subsequent interstitial deposition of fibrotic material. Fibrosis, attributed to excessive deposition of ECM proteins, is a ubiquitous tissue response to chronic inflammation [51]. In this study, ECM gene expression was altered by HFD in a time-dependent manner. Furthermore, there was evidence of accumulation of fibrillar collagens, interspersed in between adipocytes in the visceral WAT of HFD-fed mice. Lumican (*Lum*) has a stimulatory effect on the epithelial-mesenchymal transition state of fibrosis, while decorin (*Dcn*) has an inhibitory role in transforming growth factor β (TGF- β)-induced fibrosis [52,53]. *Lum* mRNA

levels were up-regulated, while *Dcn* mRNA levels were down-regulated at an early stage during the development of diet-induced obesity. *Tgfb1* mRNA levels were also up-regulated in the visceral WAT of HFD-fed mice. The up-regulation of *Lum* and *Tgfb1* mRNA levels and the down-regulation of *Dcn* mRNA levels were consistent with the observed development of fibrosis in visceral WAT of HFD-fed mice. Many metalloproteases (MMPs) were also altered at the mRNA level by HFD. *Mmp2*, *3*, *Mmp12* and *Mmp13* are reportedly positively correlated with the degree of obesity and were up-regulated in the visceral WAT of HFD fed mice [54]. Interestingly, our



detailed time-course microarray analysis over 24 weeks of HFD feeding showed that collagen mRNA levels were up-regulated shortly after the initiation of HFD feeding and then down-regulated by an unknown mechanism after 2 weeks. We suggest that up-regulation of collagen mRNA levels by HFD during the early stages of diet-induced obesity may trigger the accumulation of ECM components that ultimately leads to fibrosis which was evident in the later stages of diet-induced obesity.

The proteolytic and destructive properties of the lysosomal cathepsins play a role in degenerative, as well as chronic inflammatory diseases. Furthermore, proteases of the cathepsin family are involved in the remodelling of extracellular matrix (ECM) proteins [55]. This study indicates that members of the cathepsin family, including *Ctsb*, *Ctsk*, *Ctst*, *Ctss* and *Ctsz* as well as *Ctsa* may contribute to the pro-inflammatory and extracellular matrix remodelling in visceral WAT depots during the

development of diet-induced obesity. Destruction of elastin-rich tissues during the inflammatory response is associated with local accumulation of macrophages that contain high levels of elastinolytic enzymes, such as Ctsb and Ctsl [56]. Ctss, known to degrade several components of the ECM, is also now well established as a novel biomarker of adiposity [57]. The function of cathepsin A in obesity is relatively unknown, however increases in expression of cathepsin B, D, K, S, and Z have previously been observed in obesity [58].

It is possible that some of differences in this study compared to others could be due to the lower fat content and different diet composition. Especially, while other studies used 60 kcal% fat diet consisting of soybean oil and lard, we used 39.2 kcal% from fat diet consisting of corn oil and lard. One example, several previous studies have found that hypoxia has been observed in obese adipose tissue and HIF1 α (hypoxia-inducible factor 1 α), a master signal mediator of hypoxia response, is significantly elevated in obese adipose tissue [4,59,60]. We also found that HIF1 α gene expression tended to be higher (fold change 1.36 at HFD vs ND) at 24 week of high-fat feeding (about 39.2 kcal% from fat), but there was no statistically significant difference between the groups. However, in another experiment of ours that used commercial high fat diet with higher fat content and different fat composition (Research diets, Product No. D12451, 45 kcal% fat, 2.9% soybean oil and 20.7% lard), high fat feeding for 16 weeks significantly up-regulated HIF1 α expression in adipose tissue of C57BL/6 J mice (Y. J. Kim, M. S. Choi, U. J. Jung, unpublished paper).

Conclusions

The present findings indicate excessive fat accumulation occurs in visceral WAT depots (epididymal, perirenal, retroperitoneum, mesentery) early in response to HFD feeding. Pro- and anti-inflammatory adipokines including leptin and adiponectin are differentially expressed in visceral WAT and are closely associated with plasma adipokine levels during the development of diet-induced obesity. However, much wider changes occur in the immune transcriptome of visceral WAT depots, which are early and sustained. Up-regulated inflammatory genes include pro-inflammatory cytokines, chemokines and their upstream signalling pathway genes. Early changes also occur in fibrosis-related genes, extracellular matrix (ECM), collagen and cathepsin genes, but fibrosis appears in the later stages of diet-induced obesity. New therapeutic approaches targeting visceral adipose tissue genes associated with inflammation and fibrosis which are altered early during adipose tissue accumulation may help ameliorate the deleterious effects of diet-induced obesity.

Additional files

Additional file 1: Table S1. Composition of experimental diets. ¹AIN-76 mineral mixture (grams/kg): calcium phosphate 500, sodium chloride 74, potassium citrate 2220, potassium sulfate 52, magnesium oxide 24, magnesium carbonate 3.5, ferric citrate 6, zinc carbonate 1.6, cupric carbonate 0.3, potassium iodate 0.01, sodium selenite 0.01, chromium potassium sulfate 0.55, sucrose 118.03. ²AIN-76 vitamin mixture (grams/kg): thiamin HCL 0.6, riboflavin 0.6, pyridoxin HCL 0.7, niacin 3, calcium pantothenate 1.6, folic acid 0.2, biotin 0.02, vit B₁₂ 1, vit A(500,000U/gm) 0.8, vit D₃(400,000U/gm) 0.25, vit E acetate(500 U/gm) 10, menadione sodium bisulfite 0.08, sucrose 981.15, ND: normal diet (AIN-76), HFD: high-fat diet (20% fat, 1% cholesterol)

Additional file 2: Table S2. Primer sequences used for RT-qPCR validation of microarray data.

Additional file 3: Table S3. Body weight gain, food intake and food efficiency in C57BL/6 J mice fed the different diets for 24 weeks. Data shown as means \pm S.D. Values are significantly different from the ND group according to Student's t-test: *p < 0.05, **p < 0.01, ***p < 0.001. ND: normal diet (AIN-76), HFD: high-fat diet (20% fat, 1% cholesterol), FER: food efficiency ratio.

Additional file 4: Table S4. List of high-fat diet responding gene transcripts in epididymal WAT that simultaneously associated with advancing age in ND and HFD groups for 24 weeks.

Abbreviations

DEGs: Differentially expressed genes; ND: Normal diet; HFD: High-fat diet; WAT: White adipose tissue; ECM: Extracellular matrix.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

EYK performed experiments and wrote the manuscript. SKS and YYC carried out the animal experiments. UJJ, EK and RAM participated in writing the manuscript. TP, JHYP, JWY and YBP analyzed the data. EYK and MSC designed the study. All authors read and approved the final manuscript.

Acknowledgements

This research was supported by the SRC program (Center for Food & Nutritional Genomics: grant number 2012-0000644) of the National Research Foundation (NRF) of Korea funded by the Ministry of Education, Science and Technology.

Author details

¹Center for Food and Nutritional Genomics Research, Kyungpook National University, Daegu, Republic of Korea. ²Department of Food Science and Nutrition, Catholic University of Daegu, Gyeongsan, Republic of Korea. ³Department of Food and Nutrition, Yonsei University, Seoul, Republic of Korea. ⁴Department of Food Science and Nutrition, College of Natural Sciences, Hallym University, Chuncheon, Republic of Korea. ⁵Department of Biotechnology, Daegu University, Kyungsan, Kyungbuk, Republic of Korea. ⁶School of Life Science & Biotechnology, Kyungpook National University, Daegu, Republic of Korea. ⁷Department of Food Science and Nutrition, Kyungpook National University, 1370 Sank-Yuk Dong Puk-Ku, Daegu 702-701, Korea.

Received: 5 April 2012 Accepted: 24 August 2012

Published: 4 September 2012

References

1. Hotamisligil GS: Inflammation and metabolic disorders. *Nature* 2006, **444**(7121):860-867.
2. Rosen ED, Spiegelman BM: Adipocytes as regulators of energy balance and glucose homeostasis. *Nature* 2006, **444**(7121):847-853.
3. Maury E, Brichard SM: Adipokine dysregulation, adipose tissue inflammation and metabolic syndrome. *Mol Cell Endocrinol* 2010, **314**(1):1-16.

4. Hosogai N, Fukuhara A, Oshima K, Miyata Y, Tanaka S, Segawa K, Furukawa S, Tochino Y, Komuro R, Matsuda M, Shimomura I: **Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation.** *Diabetes* 2007, **56**(4):901–911.
5. Schönherr E, Hausser HJ: **Extracellular matrix and cytokines: a functional unit.** *Dev Immunol* 2000, **7**(2–4):89–101.
6. Okuda S, Languino LR, Ruoslahti E, Border WA: **Elevated expression of transforming growth factor-beta and proteoglycan production in experimental glomerulonephritis. Possible role in expansion of the mesangial extracellular matrix.** *J Clin Invest* 1990, **86**(2):453–462.
7. Border WA, Noble NA: **Transforming growth factor-beta in glomerular injury.** *Exp Nephrol* 1994, **2**(1):13–17.
8. Tsochatzis EA, Papatheodoridis GV, Archimandritis AJ: **Adipokines in nonalcoholic steatohepatitis: from pathogenesis to implications in diagnosis and therapy.** *Mediators Inflamm* 2009, **831670**:1–9. Published online.
9. Schram K, Sweeney G: **Implications of myocardial matrix remodeling by adipokines in obesity-related heart failure.** *Trends Cardiovasc Med* 2008, **18**(6):199–205.
10. Kleemann R, van Erk M, Verschuren L, van den Hoek AM, Koek M, Wielinga PY, Jie A, Pellis L, Bobeldijk-Pastorova I, Kelder T, Toet K, Wopereis S, Cnubben N, Evelo C, van Ommen B, Kooistra T: **Time-resolved and tissue-specific systems analysis of the pathogenesis of insulin resistance.** *PLoS One* 2010, **5**(1):e8817.
11. Khan T, Muise ES, Iyengar P, Wang ZV, Chandalia M, Abate N, Zhang BB, Bonaldo P, Chua S, Scherer PE: **Metabolic dysregulation and adipose tissue fibrosis: role of collagen VI.** *Mol Cell Biol* 2009, **29**(6):1575–1591.
12. Wang B, Wood IS, Trayhurn P: **Dysregulation of the expression and secretion of inflammation-related adipokines by hypoxia in human adipocytes.** *Pflugers Arch* 2007, **455**(3):479–492.
13. Fitzgibbons TP, Kogan S, Aouadi M, Hendricks GM, Straubhaar J, Czech MP: **Similarity of mouse perivascular and brown adipose tissues and their resistance to diet-induced inflammation.** *Am J Physiol Heart Circ Physiol* 2011, **301**(4):H1425–H1437.
14. MacLaren RE, Cui W, Lu H, Simard S, Cianflone K: **Association of adipocyte genes with ASP expression: a microarray analysis of subcutaneous and omental adipose tissue in morbidly obese subjects.** *BMC Med Genomics* 2010, **3**:3.
15. Halberg N, Khan T, Trujillo ME, Wernstedt-Asterholm I, Attie AD, Sherwani S, Wang ZV, Landskroner-Eiger S, Dineen S, Magalang UJ, Brekken RA, Scherer PE: **Hypoxia-inducible factor 1alpha induces fibrosis and insulin resistance in white adipose tissue.** *Mol Cell Biol* 2009, **29**(16):4467–4483.
16. Do GM, Kwon EY, Kim E, Kim HS, Choi MS: **Hepatic transcription response to high-fat treatment in mice: microarray comparison of individual vs. pooled RNA samples.** *Biotechnol J* 2010, **5**:970–973.
17. *In Bioinformatics and Computational Biology Solutions using R and Bioconductor.* Edited by Gentleman R, Carey V, Huber W, Irizarry R, Dudoit S. New York: Springer; 2005.
18. Schmittgen TD, Livak KJ: **Analyzing real-time PCR data by the comparative C(T) method.** *Nat Protoc* 2008, **3**:1101–1108.
19. Huang DW, Sherman BT, Lempicki RA: **Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources.** *Nat Protoc* 2009, **4**:44–57.
20. Do GM, Oh HY, Kwon EY, Cho YY, Shin SK, Park HJ, Jeon SM, Kim E, Hur CG, Park TS, Sung MK, McGregor RA, Choi MS: **Long-term adaptation of global transcription and metabolism in the liver of high-fat diet-fed C57BL/6 J mice.** *Mol Nutr Food Res* 2011, **55**(Suppl 2):S173–S185.
21. Divoux A, Clément K: **Architecture and the extracellular matrix: the still unappreciated components of the adipose tissue.** *Obes Rev* 2011, **12**(5): e494–e503.
22. Zhang F, Basinski MB, Beals JM, Briggs SL, Churgay LM, Clawson DK, DiMarchi RD, Furman TC, Hale JE, Hsiung HM, Schoner BE, Smith DP, Zhang XY, Wery JP, Schevitz RW: **Crystal structure of the obese protein leptin-E100.** *Nature* 1997, **387**(6629):206–209.
23. Tartaglia LA, Dembski M, Weng X, Deng N, Culpepper J, Devos R, Richards GJ, Campfield LA, Clark FT, Deeds J, Muir C, Sanker S, Moriarty A, Moore KJ, Smutko JS, Mays GG, Wool EA, Monroe CA, Tepper RI: **Identification and expression cloning of a leptin receptor, OB-R.** *Cell* 1995, **83**(7):1263–1271.
24. Zarkesh-Esfahani H, Pockley G, Metcalfe RA, Bidlingmaier M, Wu Z, Ajami A, Weetman AP, Strasburger CJ, Ross RJ: **High-dose leptin activates human leukocytes via receptor expression on monocytes.** *J Immunol* 2001, **167**(8):4593–4599.
25. Loffreda S, Yang SQ, Lin HZ, Karp CL, Brengman ML, Wang DJ, Klein AS, Bulkley GB, Bao C, Noble PW, Lane MD, Diehl AM: **Leptin regulates proinflammatory immune responses.** *FASEB J* 1998, **12**(1):57–65.
26. Oka T, Nishimura Y, Zang L, Hirano M, Shimada Y, Wang Z, Umemoto N, Kuroyanagi J, Nishimura N, Tanaka T: **Diet-induced obesity in zebrafish shares common pathophysiological pathways with mammalian obesity.** *BMC Physiol* 2010, **21**(10):21.
27. Steppan CM, Bailey ST, Bhat S, Brown EJ, Banerjee RR, Wright CM, Patel HR, Ahima RS, Lazar MA: **The hormone resistin links obesity to diabetes.** *Nature* 2001, **409**(6818):307–312.
28. Steppan CM, Lazar MA: **The current biology of resistin.** *J Intern Med* 2004, **255**(4):439–447.
29. Jackson MB, Osei SY, Ahima RS: **The endocrine role of adipose tissue: focus on adiponectin and resistin.** *Curr Opin Endocrin Diab* 2005, **12**:163–170.
30. Kawanami D, Maemura K, Takeda N, Harada T, Nojiri T, Imai Y, Manabe I, Utsunomiya K, Nagai R: **Direct reciprocal effects of resistin and adiponectin on vascular endothelial cells: a new insight into adipocytokine-endothelial cell interactions.** *Biochem Biophys Res Commun* 2004, **314**(2):415–419.
31. Kadowaki S, Li SH, Wang CH, Fedak PW, Li RK, Weisel RD, Mickle DA: **Resistin promotes endothelial cell activation: further evidence of adipokine-endothelial interaction.** *Circulation* 2003, **108**(6):736–740.
32. Kadowaki T, Yamauchi T, Kubota N, Hara K, Ueki K, Tobe K: **Adiponectin and adiponectin receptors in insulin resistance, diabetes, and the metabolic syndrome.** *J Clin Invest* 2006, **116**(7):1784–1792.
33. Bolduc C, Yoshioka M, St-Amand J: **Acute molecular mechanisms responsive to feeding and meal constitution in mesenteric adipose tissue.** *Obesity (Silver Spring)* 2010, **18**(2):410–413.
34. Sumiyoshi M, Sakanaka M, Kimura Y: **Chronic intake of high-fat and high-sucrose diets differentially affects glucose intolerance in mice.** *J Nutr* 2006, **136**(3):582–587.
35. Park SY, Cho YR, Kim HJ, Higashimori T, Danton C, Lee MK, Dey A, Rothermel B, Kim YB, Kalinowski A, Sakanaka M, Russell KS, Kim JK: **Unraveling the temporal pattern of diet-induced insulin resistance in individual organs and cardiac dysfunction in C57BL/6 mice.** *Diabetes* 2005, **54**(12):3530–3540.
36. Flier JS, Cook KS, Usher P, Spiegelman BM: **Severely impaired adipin expression in genetic and acquired obesity.** *Science* 1987, **237**(4813):405–8.
37. Maslowska M, Vu H, Phelis S, Sniderman AD, Rhode BM, Blank D, Cianflone K: **Plasma acylation stimulating protein, adipin and lipids in non-obese and obese populations.** *Eur J Clin Invest* 1999, **29**(8):679–686.
38. Napolitano A, Lowell BB, Damm D, Leibel RL, Ravussin E, Jimerson DC, Lesem MD, Van Dyke DC, Daly PA, Chatis P: **Concentrations of adipin in blood and rates of adipin secretion by adipose tissue in humans with normal, elevated and diminished adipose tissue mass.** *Int J Obes Relat Metab Disord* 1994, **18**(4):213–218.
39. Cianflone K, Xia Z, Chen LY: **Critical review of acylation-stimulating protein physiology in humans and rodents.** *Biochim Biophys Acta* 2003, **1609**(2):127–143.
40. Caesar R, Manieri M, Kelder T, Boeschoten M, Evelo C, Müller M, Kooistra T, Cinti S, Kleemann R, Drevon CA: **A combined transcriptomics and lipidomics analysis of subcutaneous, epididymal and mesenteric adipose tissue reveals marked functional differences.** *PLoS One* 2010, **5**(7):e11525.
41. Morton NM, Nelson YB, Michailidou Z, Di Rollo EM, Ramage L, Hadoke PW, Seckl JR, Bunger L, Horvat S, Kenyon CJ, Dunbar DR: **A stratified transcriptomics analysis of polygenic fat and lean mouse adipose tissues identifies novel candidate obesity genes.** *PLoS One* 2011, **6**(9):e23944.
42. Lin Y, Lee H, Berg AH, Lisanti MP, Shapiro L, Scherer PE: **The lipopolysaccharide-activated toll-like receptor (TLR)-4 induces synthesis of the closely related receptor TLR-2 in adipocytes.** *J Biol Chem* 2000, **275**(32):24255–24263.
43. Batra A, Pietsch J, Fedke I, Glauben R, Okur B, Stroh T, Zeitz M, Siegmund B: **Leptin-dependent toll-like receptor expression and responsiveness in preadipocytes and adipocytes.** *Am J Pathol* 2007, **170**(6):1931–1941.
44. Takaoka A, Yanai H, Kondo S, Duncan G, Negishi H, Mizutani T, Kano S, Honda K, Ohba Y, Mak TW, Taniguchi T: **Integral role of IRF-5 in the gene induction programme activated by Toll-like receptors.** *Nature* 2005, **434**(7030):243–249.
45. Beutler B: **Tlr4: central component of the sole mammalian LPS sensor.** *Curr Opin Immunol* 2000, **12**(1):20–26.

46. Miyake K: **Roles for accessory molecules in microbial recognition by Toll-like receptors.** *J Endotoxin Res* 2006, **12**(4):195–204.
47. Akashi-Takamura S, Miyake K: **TLR accessory molecules.** *Curr Opin Immunol* 2008, **20**(4):420–425.
48. Ogata H, Su I, Miyake K, Nagai Y, Akashi S, Mecklenbräuker I, Rajewsky K, Kimoto M, Tarakhovskiy A: **The toll-like receptor protein RP105 regulates lipopolysaccharide signaling in B cells.** *J Exp Med* 2000, **192**(1):23–29.
49. Pierleoni C, Verdenelli F, Castellucci M, Cinti S: **Fibronectins and basal lamina molecules expression in human subcutaneous white adipose tissue.** *Eur J Histochem* 1998, **42**(3):183–188.
50. Lijnen HR: **Angiogenesis and obesity.** *Cardiovasc Res* 2008, **78**(2):286–293.
51. Wynn TA: **Common and unique mechanisms regulate fibrosis in various fibroproliferative diseases.** *J Clin Invest* 2007, **117**(3):524–529.
52. Kolb M, Margetts PJ, Sime PJ, Gauldie J: **Proteoglycans decorin and biglycan differentially modulate TGF-beta-mediated fibrotic responses in the lung.** *Am J Physiol Lung Cell Mol Physiol* 2001, **280**(6):L1327–L1334.
53. Rada JA, Cornuet PK, Hassell JR: **Regulation of corneal collagen fibrillogenesis in vitro by corneal proteoglycan (lumican and decorin) core proteins.** *Exp Eye Res* 1993, **56**(6):635–648.
54. Chavey C, Mari B, Monthouel MN, Bonnafous S, Anglard P, Van Obberghen E, Tartare-Deckert S: **Matrix metalloproteinases are differentially expressed in adipose tissue during obesity and modulate adipocyte differentiation.** *J Biol Chem* 2003, **278**(14):11888–11896. Epub.
55. McGrath ME: **The lysosomal cysteine proteases.** *Annu Rev Biophys Biomol Struct* 1999, **28**:181–204.
56. Fiebiger E, Maehr R, Villadangos J, Weber E, Erickson A, Bikoff E, Ploegh HL, Lennon-Duménil AM: **Invariant chain controls the activity of extracellular cathepsin L.** *J Exp Med* 2002, **196**(9):1263–1269.
57. Taleb S, Lacasa D, Bastard JP, Poitou C, Cancellato R, Pelloux V, Viguierie N, Benis A, Zucker JD, Bouillot JL, Coussieu C, Basdevant A, Langin D, Clement K: **Cathepsin S, a novel biomarker of adiposity: relevance to atherogenesis.** *FASEB J* 2005, **19**(11):1540–1542.
58. Nadler ST, Stoehr JP, Schueler KL, Tanimoto G, Yandell BS, Attie AD: **The expression of adipogenic genes is decreased in obesity and diabetes mellitus.** *Proc Natl Acad Sci USA* 2000, **97**(21):11371–11376.
59. Rausch ME, Weisberg S, Vardhana P, Tortoriello DV: **Obesity in C57BL/6 J mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration.** *Int J Obes (Lond)* 2008, **32**(3):451–463.
60. Ye J, Gao Z, Yin J, He Q: **Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary obese mice.** *Am J Physiol Endocrinol Metab* 2007, **293**(4):E1118–E1128.

doi:10.1186/1471-2164-13-450

Cite this article as: Kwon *et al.*: Time-course microarrays reveal early activation of the immune transcriptome and adipokine dysregulation leads to fibrosis in visceral adipose depots during diet-induced obesity. *BMC Genomics* 2012 **13**:450.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

